## MEDICATIONS TO AVOID

## 7 TO 10 DAYS BEFORE YOUR SURGERY (UNLESS OTHERWISE INSTRUCTED)

This is a list of frequently used blood thinning medications. However, your medication may not be included here. Please call our office if you are unsure whether to stop the medicine before surgery.

\*Your Primary Care Physician (PCP) and or Cardiologist must fax over a written statement that is it okay to stop major blood thinning medications prior to surgery. Please have them fax this permission statement to our office at 540-301-2579.

ASPIRIN 81 MG
ASPIRIN 325 MG
ALL BAYER ASPIRIN
CILOSTAZOL (PLETAL)
CLOPIDOGREL/PLAVIX (AS

DIRECTED)

COUMADIN /WARDARIN (AS

DIRECTED)

AGGRENOX (AS DIRECTED)

**EXCEDRIN** 

**ELIQUIS (APIXABAN)** 

**MOTRIN** 

PLETAL (CLLOSTAZOL) STOP

FOR 4 DAYS)

PRADAXA / DABIGATRAN PLETAL / CILOSTAZOL ST JOSEPHS ASPIRIN (FOR

CHILDREN)

XARELTO / RIVAROXABAN

ACCUTANE ADVIL

**ALKA-SELTZER TABLETS** 

ANAPROX APC TABLETS

**BC POWDER AND TABLETS** 

BOXTRA
CELEBREX
CELECOXIB

CHINESE HERBALS

COMBUNOX

**CONGESPIRIN** 

COPE CORLCIDIN DARVON

DAYPRO (OXAPROZIN) DICLOFENAC SODLUM

VOLTAREN DIPYRIDAMOLE ECOTRIN

EQUAGISIC FELDENE

FENOPROFEN (NALFON)

FISH OLL

FLAX SEED FLURBIPROFEN

**GINGER** 

GINGKO BILOBA GLUCOSAMINE

GOODY'S HEADACHE POWDERS

IBUPROFIN

I-CAPS INDOCIN

INDOMETHACIN KETOPROFEN

LODINE (ETODOLAC)

LOVAZA

MEFANAMIC ACID (PONSTEL)

MELOXICAM

MIDOL

MOMENTUM MUSCULAR

MONO-GESIC

NAPROSYN/NAPROXEN NORGESIC/NORGESIC FORTE

NORWICH ASPIRIN OXAPROZIN (DAYPRO)

PONSTEL (MEFENAMIC ACID)

PRADAXA-PRODUCTS

QUIET WORLD SLEEPING ALD

**RELAFEN** 

(RESERVATROL (STOP 2 WEEKS BEFORE AND 2 WEEKS AFTER

SURGERY)

ROBAXISAL TABLETS SALSALATE

ST JOHN'S WORT

STERO-DARVON WITH ASPIRIN

TORADOL (KETEROLAC)
TRENTAL (AS DIRECTED)

VALDECOCOXIB
VIOXX (AS DIRECTED)

VITAMIN E

**VOLTAREN (DICLOFENAC** 

SODIUM) VICOPROFEN

WHITE WILLOW- BARK